Write your name here		
Surname	Other nam	es
Pearson Edexcel Level 1/Level 2 GCSE (9–1)	Centre Number	Candidate Number
Physics Paper 1	Fo	undation Tier
Sample Assessment Material for first to Time: 1 hour 45 minutes	Paper Reference 1PH0/1F	
You must have: Calculator, ruler, protractor		Total Marks

## **Instructions**

- Use **black** ink or ball-point pen.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer all questions.
- Answer the questions in the spaces provided
  - there may be more space than you need.
- Calculators may be used.
- Any diagrams may NOT be accurately drawn, unless otherwise indicated.
- You must **show all your working out** with **your answer clearly identified** at the **end of your solution**.

#### Information

- The total mark for this paper is 100.
- The marks for each question are shown in brackets
   use this as a guide as to how much time to spend on each question.
- In questions marked with an asterisk (\*), marks will be awarded for your ability to structure your answer logically showing how the points that you make are related or follow on from each other where appropriate.

#### **Advice**

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ▶

**PEARSON** 


S50049A
©2016 Pearson Education Ltd.



## Answer ALL questions. Write your answers in the spaces provided.

Some questions must be answered with a cross in a box  $\boxtimes$ . If you change your mind about an answer, put a line through the box  $\boxtimes$  and then mark your new answer with a cross  $\boxtimes$ .

1 (a) Figure 1 shows a solar-powered charger for a mobile phone.



The screen of the solar cell takes in energy from the Sun.

(i) State how energy gets from the Sun to the screen.

(1)

(ii) State how energy is stored in the charger.

b) Mobile phones emit microwaves.  Microwave ovens emit microwaves.		(Total for Question 1 = 7 marks)
energy =b) Mobile phones emit microwaves.  Microwave ovens emit microwaves.  Explain why a mobile phone does not have the same heating effect as a microwave oven.		
energy =b) Mobile phones emit microwaves.  Microwave ovens emit microwaves.  Explain why a mobile phone does not have the same heating effect as a microwave oven.		
energy =b) Mobile phones emit microwaves.  Microwave ovens emit microwaves.  Explain why a mobile phone does not have the same heating effect as a microwave oven.		
energy =b) Mobile phones emit microwaves.		
energy =		Microwave ovens emit microwaves.
	b)	Mobile phones emit microwaves.
		energy =

**2** (a) A student is standing 600 m from a firework display.

A firework explodes with a loud bang, and a flash of light is seen.

Describe how a student can measure the time it takes for the sound wave from the loud bang to travel 600 m.

(2)

(b) Figure 2 shows a water wave.




Figure 2

(i) What is the amplitude of this wave?

(1)

- **A** 2.8 cm
- **■ B** 5.6 cm

(ii) What is the wavelength of this wave?

- A 2.8 cm
- B 7.5 cm
- **∑** 15 cm

(	(c)	Water	waves	are	transverse	waves
•	( )	vvatei	waves	are	Hallsveise	waves

(i) Give **one** other example of a transverse wave.

(1)

(ii) Give **one** example of a longitudinal wave.

(1)

(d) An earthquake causes a sea wave.

This sea wave travels 26400 m in two minutes.

Calculate the speed of the wave.

Use the equation

$$wave speed = \frac{distance}{time}$$

(3)

(Total for Question 2 = 9 marks)

- **3** (a) Figure 3 shows the structure of an oxygen-14 atom.
  - (i) Complete the four labels on Figure 3.






Figure 3

(ii) Which of these particles has a negative charge?

(1)

- A alpha particle
- B electron
- **D** nucleus
- (iii) State the overall charge of the oxygen-14 atom.

		(Total for Question 3 = 9 mar	ks)					
	(iii) 	Give a reason why the answer to (ii) is only an estimate.	per minute					
		Estimate the reading on the counter tube.	(1)					
	(ii)	The teacher puts a thick sheet of aluminium between the source of beta radiation and the Geiger-Müller tube.	per minute					
		Calculate how much radiation detected by the Geiger-Müller tube comes from the source of beta radiation.	(1)					
		tube.  The reading on the counter tube is now 468 counts per minute.						
		The teacher puts a source of beta radiation 15 cm in front of the same Geiger-M	üller					
		e reading on the counter tube is 34 counts per minute.						
(b)	Αt	eacher uses a Geiger-Müller tube and a counter to measure background radiatio	n.					

## **BLANK PAGE**

**4** A student investigates how light behaves as it leaves a clear plastic block.

Figure 4 shows some of her equipment and the path of a ray of light through the block.




Figure 4

(a) Describe how the student can make sure the light does not change direction as it enters the block.


(2)

(b) She varies the angle of incidence inside the block and records the angle of refraction. Figure 5 shows her results.


angle of incidence (°)	angle of refraction (°)
5	7
15	22
30	46
40	69
42	76

Figure 5

(i) Plot the points on the grid below.

(2)

(ii) Draw the best fit smooth curve through the points.



(Total for Question 4 =	
(c) Describe what happens to the ray of light when the angle of incidence is increased beyond the critical angle.	(2)
angle of incidence =	·
(iii) Estimate the angle of incidence which gives an angle of refraction of 90°.	(2)

**5** (a) A car driver sees a rabbit on the road.

The driver makes an emergency stop after he sees the rabbit.

Figure 6 shows the speed of the car from the time the driver sees the rabbit until the car stops.




Figure 6

(i) The distance travelled by the car from the time the driver first sees the rabbit to when car starts to slow down is the

(1)

- A average distance
- B braking distance
- C stopping distance
- ☑ D thinking distance
- (ii) Calculate the distance that the car travels in the first 0.5 seconds.

(3)

distance = ..... m

(iii) Which equation relates acceleration to change in velocity and time?

(1)

- $\triangle$  **A**  $a = \frac{(v-u)}{t}$
- $\blacksquare$  **B**  $a = \frac{t}{(v-u)}$
- $\square$  **C** a = t(v u)
- $\square$  **D**  $a = v \frac{u}{t}$
- (iv) Calculate the deceleration of the car.

(3)

deceleration = ..... m/s<sup>2</sup>

	(Total for Question 5 = 11 marks)
3	
2	
1	
	Give <b>three</b> ways the students could improve their experimental procedure. (3)
	'My estimate for the time taken for the car to pass between the two lamp posts = $3'$
	He records:
	Bob starts to count when a car passes the first lamp post. He stops counting when he thinks it has passed the second lamp post.
	'Distance between lamp posts = 20 paces'
	She records:
	Alice paces out the distance between two lamp posts.
(I	b) Two students, Alice and Bob, carry out an experiment to measure the speed of cars.

**6** (a) A student investigates how the surface of an object affects the radiation it emits.

Figure 7 shows the equipment he uses.




Figure 7

The cube has four different surfaces.


He fills the cube with boiling water so that the temperature of each surface is the same.

He uses the radiation sensor to measure the radiation emitted from each surface.

(i) His readings are shown.

Draw a line from each surface colour to its correct meter reading. One has been done for you.

(2)



	(ii)	The temperature of each surface is the same.  Give a reason why the radiation sensor gives a different reading for each surface.	ce. (1)
(b)	(i)	What do all waves in the electromagnetic spectrum have in common?	(1)
X	Α	the same frequency in a vacuum	
×	В	the same speed in a vacuum	
X	C	the same colour in a vacuum	
X	D	the same amplitude in a vacuum	
	(ii)	Blue light has a wavelength of 470 nm and a frequency of $6.30\times10^{14}\text{Hz}$ Calculate the velocity of blue light.	(2)

velocity = ..... m/s

(c) All objects emit electromagnetic radiation.

The intensity and wavelength of the emitted radiation vary with the temperature of the object.

Figure 8 shows this variation for an object at two different temperatures.

The visible region of the electromagnetic spectrum is also shown.




Figure 8

(i) In which part of the electromagnetic spectrum is the peak of the low temperature curve?

(1)

- 🛚 🗛 gamma
- B infrared
- C radio
- D ultra violet
- (ii) Describe how intensity of the emitted radiation changes with temperature.

(2)

(Total for Question 6 = 9 marks)

7	(a)	Scientists no longer accept the geocentric model of the universe but it was the accepted theory for hundreds of years.	
		Explain why the evidence available at the time supported the geocentric model.	
			(3)
	(b)	The Big Bang theory and the Steady State theory are two theories about the origin of the universe.	
		The discovery of CMB led scientists to accept only one of the theories.	
		Explain why redshift supports both theories but CMB supports only one of them.	
			(3)
•••••			
•••••	•••••		

(c)	(i)	A s	tar with a mass very much larger than the Sun	(1)
	X	A	has a longer main sequence than the Sun and ends as a white dwarf	
	X	В	has a longer main sequence than the Sun and ends as a black hole	
	X	C	has a shorter main sequence than the Sun and ends as a white dwarf	
	X	D	has a shorter main sequence than the Sun and ends as a black hole	

(ii) Which row has two correct statements about black holes?

		the gravitational field of a black hole	a black hole is formed when
X	A	allows only electromagnetic radiation to escape	a nebula collapses
×	В	allows nothing to escape	a very large star collapses
X	c	allows nothing to escape	a nebula collapses
X	D	allows only electromagnetic radiation to escape	a very large star collapses

(d) Figure 9 shows some lines in the absorption spectra from four different galaxies (A, B, C, and D) and from a laboratory source.

All the spectra are aligned and to the same scale.




Figure 9

Explain, using Figure 9, which galaxy is furthest away from us.

(3)

(Total for Question 7 = 11 marks)

**8** Figure 10 shows two students investigating reaction times.

Student B supports his left hand on a desk.

Student A holds a ruler so that the bottom end of the ruler is between the finger and thumb of student B.

When student A releases the ruler, student B catches the ruler as quickly as he can with his left hand.

The investigation is repeated with the right hand of student B.




Figure 10

(a)	Give a reason why it is better to have the 0 cm mark at the bottom of the rule
	rather than at the top.

(1)

(b) Give a reason why two students are needed for this investigation.

(c) The students took five results for the left hand and five results for the right hand.

Figure 11 shows their results.

which			distance dr	opped (cm)		
hand	trial 1	trial 2	trial 3	trial 4	trial 5	average
left	10.1	25.5	18.4	14.6	11.7	14
right	17.5	16.1	19.4	18.6	20.2	

Figure 11

(i) Calculate the average distance dropped for the right hand. Give your answer correct to 2 significant figures.

(2)

(ii) Calculate the average time for the left hand.

Use the equation

$$time^2 = \frac{distance}{500}$$

(2)

(d) Explain whether any of t	e readings are anomalous.	(2)
(e) Give <b>two</b> ways that the ignoring anomalous res	udents can improve the quality of their data, other tha	
		(2)
(f) Describe how the stude reaction time changes v	s could develop their investigation to investigate how h another variable.	(2)
	(Total for Question 8 = 12 n	narks)

- **9** A car accelerates at a constant rate of 1.83 m/s² along a flat straight road.
  - (a) The force acting on the car is 1.870 kN.

Calculate the mass of the car.

Give your answer to three significant figures.

(3)

(b) The car accelerates from rest for 16 s.

Calculate the speed of the car after 16 s.

(3)

\*(c) Figure 12 is a speed-time graph for a different car moving on a horizontal road.




Figure 12

Describe the energy transfers taking place during the movement of the car.

You should refer to energy stores as well as transfers between energy stores for all three sections of the graph.

(Total for Question 9 = 12 marks)

(6)

10	(a)	Americium-241 is a radioactive isotope that emits alpha particles.  Americium-241 is used in smoke alarms.	
		Give a reason why it is safe to use americium-241 in smoke alarms.	(1)
	(b)	lodine-131 is a radioactive isotope with a half-life of 8 days.	
		The activity of a sample of iodine-131 is 480 Bq.	
		Calculate the activity of the sample after 16 days.	(2)
		activity =	Bq
	(c)	A student uses 59 dice to model radioactive decay.	
		He starts by rolling all the dice at the same time.	
		He removes all the dice that show a six.	
		He then rolls the remaining dice.	
		The student repeats this process five more times.	
		State <b>two</b> improvements the student could make to his model of radioactive decay.	(2)
1			
2			

*(d) Dadioactive isotomes can be used to investigate concerned ather illnesses	
*(d) Radioactive isotopes can be used to investigate cancer and other illnesses.	
The thyroid gland in the neck absorbs most of the iodine that our bodies need.	
A person can become ill if their thyroid absorbs too little iodine.	
Explain how a radioactive isotope with suitable properties may be used to investigate the uptake of iodine by this gland.	
	(6)
(Total for Question 10 = 11 n	narks)
TOTAL FOR PAPER = 100 M	ARKC

## **Equations**

(final velocity)<sup>2</sup> – (initial velocity)<sup>2</sup> =  $2 \times acceleration \times distance$ 

$$v^2 - u^2 = 2 \times a \times x$$

energy transferred = current  $\times$  potential difference  $\times$  time

$$E = I \times V \times t$$

potential difference across primary coil  $\times$  current in primary coil = potential difference across secondary coil  $\times$  current in secondary coil

$$V_p \times I_p = V_s \times I_s$$

change in thermal energy = mass  $\times$  specific heat capacity  $\times$  change in temperature

$$\Delta Q = m \times c \times \Delta \theta$$

thermal energy for a change of state =  $mass \times specific$  latent heat

$$Q = m \times L$$

$$P_1 V_1 = P_2 V_2$$

to calculate pressure or volume for gases of fixed mass at constant temperature

energy transferred in stretching =  $0.5 \times \text{spring constant} \times (\text{extension})^2$ 

$$E = \frac{1}{2} \times k \times x^2$$

# **Paper 1 Foundation**

Question number	Answer	Mark
1(a)(i)	(Carried by) electromagnetic wave	(1)

Question number	Answer	Mark
1(a)(ii)	As chemical energy in the battery	(1)

Question number	Answer	Additional guidance	Mark
1(a)(iii)	Calculation of area (1) 7 × 11	77	
	Substitution (1) 77 × 0.12	ecf area	
	Answer (1) 9.2 (J)	award full marks for correct numerical answer without working	(3)

Question number	Answer	Additional guidance	Mark
1(b)	An explanation that combines identification – understanding (1 mark) and reasoning/justification – understanding (1 mark):  • the heating effect for the oven and the phone depends on their power (1)  • and since the power of an oven is much greater than the power of a phone, the oven produces a greater heating effect (1)	allow not the same wavelength/microwaves cover a range in wavelengths	(2)

Question number	Answer	Mark
2(a)	An answer that combines the following points of understanding to provide a logical description:	
	<ul> <li>use a stopwatch (1)</li> <li>start timing when flash is seen and stop when bang is heard (1)</li> </ul>	(2)

Question number	Answer	Mark
2(b)(i)	A	(1)

Question number	Answer	Mark
2(b)(ii)	С	(1)

Question number	Answer	Additional guidance	Mark
2(c)(i)	electromagnetic wave	allow any named e.m. wave/seismic S wave	(1)

Question number	Answer	Additional guidance	Mark
2(c)(ii)	sound wave	allow ultrasound/infrasound/ seismic P wave	(1)

Question number	Answer	Additional guidance	Mark
2(d)	two minutes = 120 s (1)  substitution (1) 26 400 ÷ 120  answer (1) 220 (m/s)	ecf unit change award full marks for correct numerical answer without working	
			(3)

Question number	Answer	Mark
3(a)(i)	One mark for each correct label (4)  electron nucleus	
		(4)

Question number	Answer	Mark
3(a)(ii)	В	(1)

Question number	Answer	Mark
3(a)(iii)	zero/0/no charge	(1)

Question number	Answer	Mark
3(b)(i)	434	(1)

Question number	Answer	Additional guidance	Mark
3(b)(ii)	34	allow 29 to 39	(1)

Question number	Answer	Additional guidance	Mark
3(b)(iii)	Radioactive decay is a random process	allow because background count changes every time	(1)

Question number	Answer	Additional guidance	Mark
4(a)	An answer that combines the following points of understanding to provide a logical description:  • shine the light along a radius (1)  • by marking it on the paper before putting the block down (1)	shine the ray at the centre of the straight edge before putting the block down	(2)

Question number	Answer	Additional guidance	Mark
4(b)(i)	all points correctly plotted to +/- half a square (2)	4 points plotted correctly (i.e. one error) (1)	
			(2)

Question number	Answer	Mark
4(b)(ii)	smooth curve through at least 3 of the points (1)	(1)

Question number	Answer	Additional guidance	Mark
4(b)(iii)	<ul> <li>continues line as far as 90° (1)</li> <li>estimate between 43° and 47° (1)</li> </ul>	award full marks for correct numerical answer without working	(2)

Question number	Answer	Mark
4(c)	An answer that provides a description by making reference to:	
	<ul><li>(all) light reflected (1)</li><li>back inside block (1)</li></ul>	(2)

Question number	Answer	Mark
5(a)(i)	D	(1)

Question number	Answer	Additional guidance	Mark
5(a)(ii)	16.0 (m/s) read from graph (1)		
	Substitution (1) (distance travelled =) 16 × 0.5 Answer (1) 8.0 (m) (1)	award full marks for correct numerical answer without working  ecf for substitution and answer using wrong speed value	
			(3)

Question number	Answer	Mark
5(a)(iii)	A	(1)

Question number	Answer	Additional guidance	Mark
5(a)(iv)	Obtain readings from graph (1)  Substitution (1) $\frac{16}{2.0}$ Answer (1)	award full marks for correct numerical answer without working	
	8.0 (m/s <sup>2</sup> )		(3)

Question number	Answer	Additional guidance	Mark
5(b)	<ul> <li>Any three improvements from:</li> <li>suitable instrument to measure distance (1)</li> <li>using a greater distance (to reduce effect of reaction times) (1)</li> <li>suitable instrument to measure time (1)</li> <li>use of one student at the {first/second} lamp post to signal when to {start/stop} timing (1)</li> </ul>	allow tape measure, trundle wheel allow stop watch/clock or timing app. on phone	(3)

two of three sets of students taking readings for the same
car (1)

Question number	Answer	Mark
6(a)(i)	All three correct (2)	
	One or two correct (1)	
	shiny black 87	
	dull black 61	
	dull silver 70	
	shiny silver 47	
		(2)

Question number	Answer	Additional guidance	Mark
6(a)(ii)	Different surfaces emit (thermal) radiation at different rates	allow reference to surfaces in question	
			(1)

Question number	Answer	Mark
6(b)(i)	В	(1)

Question number	Answer	Additional guidance	Mark
6(b)(ii)	substitution and unit conversion (1) $470 \times 10^{-9} \times 6.30 \times 10^{14}$	award full marks for correct numerical answer without working	
	answer (1) 2.96 × 10 <sup>8</sup> (m/s)	ecf unit conversion	(2)

Question number	Answer	Mark
6(c)(i)	В	(1)

Question number	Answer	Mark
6(c)(ii)	<ul> <li>An answer that combines points of interpretation/evaluation to provide a logical description:</li> <li>as temperature increases, intensity increases (1)</li> <li>as temperature increases, maximum intensity occurs at a shorter wavelength (1)</li> </ul>	(2)

Question number	Answer	Additional guidance	Mark
7(a)	An explanation that combines identification – understanding (1 mark) and reasoning/justification – understanding (2 marks):  • at the time, there was only naked eye evidence (1)  • which indicated Sun/Moon/planets appear to move across the sky (1)  • in the same direction, same motion each day (1)	allow valid alternatives, e.g. references to comets	(3)

Question number	Answer	Additional guidance	Mark
7(b)	An explanation that combines identification – understanding (1 mark) and reasoning/justification – understanding (2 marks):  • both theories predict an		
	<ul> <li>both theories predict an expanding universe and the Big Bang theory also predicts that the universe had a beginning (1)</li> <li>the red shift theory indicates that the universe is expanding so supports both theories (1)</li> <li>whereas CMB also indicates that the universe had a</li> </ul>	provided evidence that the steady state theory was incorrect	
	beginning, so supports Big Bang theory (1)		(3)

Question number	Answer	Mark
7(c)(i)	В	(1)

Question number	Answer	Mark
7(c)(ii)	В	(1)

Question number	Answer	Mark
7(d)	An explanation that combines identification via a judgement (1 mark) to reach a conclusion via justification/reasoning (2 marks):	
	<ul> <li>galaxy C is furthest away (1)</li> <li>because it has the greatest red shift (1)</li> <li>and therefore it has the greatest speed (1)</li> </ul>	(3)

Question number	Answer	Mark
8(a)	Idea of a direct reading (without calculation)	(1)

Question number	Answer	
8(b)	If student B drops the ruler, they are not really measuring their own reaction time as they know when ruler has been dropped	
		(1)

Question number	Answer	Additional guidance	Mark
8(c)(i)	calculating the mean (1) 18.36	award full marks for correct numerical answer without working	
	rounding to 2 s.f. (1) 18 (cm)		(2)

Question number	Answer	Additional guidance	Mark
8(c)(ii)	Rearrangement (1) $t = \sqrt{\frac{\text{distance}}{500}}$	award full marks for correct numerical answer without working	
	Substitution and answer (1) time = 0.17 (s)	allow answers which round to 0.17, e.g. 0.1673	(2)

Question number	Answer	Additional guidance	Mark
8(d)	An explanation that combines identification via a judgement (1 mark) to reach a conclusion via justification/reasoning (1 mark):		
	<ul> <li>25.5 is an anomalous result <ul> <li>(1)</li> <li>(because) it is much further away from the mean than the other results (1)</li> </ul> </li> </ul>	ignore 19	(2)

Question number	Answer	Mark
8(e)	<ul><li>Take more readings (1)</li><li>Idea that a third student should also measure the</li></ul>	
	reaction time (1)	(2)

Question number	Answer	Additional guidance	Mark
8(f)	An answer that combines the following points to provide a logical description of the plan/method/experiment:  • using a larger group of students/large population of students (1)  • and measure how their reaction time varies with age/height (1)	allow any suitable variable	(2)
			(2)

Question number	Answer	Additional guidance	Mark
9(a)	rearrangement (1) $m = \frac{f}{a}$ substitution and conversion (1) $m = \frac{1870}{1.83}$ answer and rounding to 3 s.f. (1) 1020 (kg)	maximum 2 marks if kN not converted to N award full marks for correct numerical answer without working	(3)

Question number	Answer	Additional guidance	Mark
9(b)	rearrangement of $\frac{(v-u)}{t} = a$ (1)		
	$v = u + at$ substitution (1) $v = 0 + 1.83 \times 16$		
	answer (1) 29.3 (m/s)	award full marks for correct numerical answer without working	(3)

Question number	Indicative content	Mark
*9(c)	Answers will be credited according to candidate's deployment of knowledge and understanding of the material in relation to the qualities and skills outlined in the generic mark scheme.	
	The indicative content below is not prescriptive and candidates are not required to include all the material which is indicated as relevant. Additional content included in the response must be scientific and relevant.	
	AO2	
	<ul> <li>fuel forms a store of chemical (potential) energy</li> <li>chemical energy is transferred to kinetic energy and thermal energy when the car moves</li> <li>kinetic energy transferred to thermal energy as the car slows down</li> </ul>	
	AO3	
	<ul> <li>during X, kinetic energy increases as the car's speed increases/car accelerates and the increase in kinetic energy is provided by the chemical energy store</li> </ul>	
	<ul> <li>during all three sections, work is done against frictional forces in the moving parts of the car and against the drag from the air</li> </ul>	
	<ul> <li>during Y, kinetic energy stays constant when the car moves at constant speed but energy is still transferred to thermal energy</li> </ul>	
	during Z, kinetic energy decreases as the car slows down	(6)

Level	Mark	Descriptor
	0	No awardable content.
1	1-2	<ul> <li>Interpretation and evaluation of the information attempted but will be limited with a focus on mainly just one variable. Demonstrates limited synthesis of understanding. (AO3)</li> <li>The description attempts to link and apply knowledge and understanding of scientific ideas, flawed or simplistic connections made between elements in the context of the question. (AO2)</li> </ul>
2	3-4	<ul> <li>Interpretation and evaluation of the information on both variables, synthesising mostly relevant understanding. (AO3)</li> <li>The description is mostly supported through linkage and application of knowledge and understanding of scientific ideas, some logical connections made between elements in the context of the question. (AO2)</li> </ul>

3	5-6	Interpretation and evaluation of the information, demonstrating throughout the skills of synthesising relevant understanding. (AO3)
		<ul> <li>The description is supported throughout by linkage and application of knowledge and understanding of scientific ideas, logical connections made between elements in the context of the question. (AO2)</li> </ul>

Question number	Answer	Additional guidance	Mark
10(a)	alpha cannot penetrate casing	alpha only travel a few cm in air	(1)

Question number	Answer	Mark
10(b)	<ul><li>evidence of division of activity by 2 (1)</li><li>120 (Bq) (1)</li></ul>	(2)

Question number	Answer	Mark
10(c)	<ul><li>increase number of starting dice (1)</li><li>do more rolls (1)</li></ul>	(2)

Question	Indicative content	Mark
number		

*10(d)	Answers will be credited according to candidate's		
(/	deployment of knowledge and understanding of the		
	material in relation to the qualities and skills outlined in		
	the generic mark scheme.		
	The indicative content below is not prescriptive and candidates are not required to include all the material which is indicated as relevant. Additional content included in the response must be scientific and relevant.		
	AO2 (6 marks)		
	use a radioactive isotope of iodine as this is taken up by the gland		
	isotope given by injection or orally		
	gland is in the neck, so cannot use an alpha emitter as		
	alpha will not exit through the skin		
	use beta or gamma emitter		
	<ul> <li>isotope has to have a short enough half-life to minimise exposure to radiation but long enough for the reading to be taken</li> </ul>		
	allow time for isotope to reach gland		
	use Geiger-Müller tube and counter to determine count		
	rate of isotope in gland		
	compare with normal count rate to determine whether		
	uptake of iodine is normal	(6)	
	aptance of loanie is normal	1 ( 0 )	

Level	Mark	Descriptor
	0	No awardable content.
1	1-2	<ul> <li>The explanation attempts to link and apply knowledge and understanding of scientific ideas, flawed or simplistic connections made between elements in the context of the question. (AO2)</li> <li>Lines of reasoning are unsupported or unclear. (AO2)</li> </ul>
2	3-4	<ul> <li>The explanation is mostly supported through linkage and application of knowledge and understanding of scientific ideas, some logical connections made between elements in the context of the question. (AO2)</li> <li>Lines of reasoning mostly supported through the application of relevant evidence. (AO2)</li> </ul>
3	5-6	<ul> <li>The explanation is supported throughout by linkage and application of knowledge and understanding of scientific ideas, logical connections made between elements in the context of the question. (AO2)</li> <li>Lines of reasoning are supported by sustained application of relevant evidence. (AO2)</li> </ul>